

Table 1

|        | Anesthetic | Medications   | Weight  | Height   | Comorbidites  |  |
|--------|------------|---|---------|----------|---|--|
| Case 1 | Spinal     | No local anesthetic noted; Fentanyl 10 mcg; Morphine 100 mcg            | 60 kg   | 134.9 cm | Achondroplasia; Thoracic rods; L2 - L3 laminectomy; Tonsil and adenoidectomy          |  |
| Case 2 | CSE        | Bupivacaine 7.5 mg; Fentanyl 10 mcg; Morphine 100 mcg; Clonidine 50 mcg | 80.1 kg | 147.3 cm | IMAGe syndrome; Pre-Eclampsia; SVT with WPW, Multiple previous abortions              |  |
| Case 3 | CSE        | Bupivacaine 12 mg; Fentanyl 25 mcg; Morphine 100 mcg                    | 67.5 kg | 153 cm   | Hereditary Multiple Exostosis   |  |
| Case 4 | CSE        | Bupivacaine 5mg; Fentanyl 10 mcg  | 42 kg   | 91.9 cm  | Osteogenesis Imperfecta type 3; Asthma; Scoliosis                                     |  |
| Case 5 | CSE        | Bupivacaine 3 mg; Fentanyl 10 mcg; Morphine 100 mcg                     | 50 kg   | 120 cm   | Achondroplasia; Central sleep apnea, Gestational diabetes, Atlanto-axial instability  |  |
| Case 6 | GA         | TIVA  | 50 kg   | 83.1 cm  | Spondyloepiphyseal dysphasia congenita; Restrictive lung disease (unable to lie flat) |  |
| Case 7 | GA         | Sevoflurane, Propofol, Remifentanil                                     | 36 kg   | 111 cm   | Osteogenesis Imperfecta; Restrictive lung disease; Difficult airway; Scoliosis        |  |