

Table 1. Unanticipated Birth Events Analyzed

Events	Subcategories Analyzed
<i>Neonatal Intensive Care Unit admission</i>	n/a
<i>Unanticipated Induction/Augmentation Indication</i>	<ul style="list-style-type: none"> a. <u>Maternal serious comorbidity</u>; Pre-eclampsia without severe features, Pre-eclampsia with severe features, Gestational Hypertension b. <u>Maternal comorbidity</u>; Diabetes, Infection, Cholestasis, c. <u>Fetal serious comorbidity</u>; IUGR, Non-reassuring fetal status d. Failure to progress
<i>Unanticipated Operative Delivery Indication</i>	<ul style="list-style-type: none"> a. Vacuum, Forceps b. Non-reassuring fetal status c. Arrest of dilation, Arrest of descent Failed Trial of labor after cesarean (TOLAC)
<i>Hemorrhage Management</i>	<ul style="list-style-type: none"> a. <u>3 or more medications</u>; 2nd dose oxytocin, methergine (methylergonovine maleate), cytotec (misoprostol), hemabate (carboprost) b. Manual removal of placenta c. <u>Operative management</u>; B-lynch, Bakri, D&C, Hysterectomy, Interventional radiology d. Blood transfusion e. ICU admission
<i>Pain with labor epidural</i>	<ul style="list-style-type: none"> a. # of times called into the room for pain b. # of times epidural replaced c. > or = 3 attempts OR > 45minutes to place epidural d. Pain score of 8, 9, or 10 with labor epidural in place
<i>Pain during cesarean delivery</i>	<ul style="list-style-type: none"> a. Epidural lidocaine bolus b. <u>Intravenous medications</u>; ketamine, narcotics, midazolam c. General anesthesia, Nitrous oxide d. Reassurance
<i>Anticipated Inductions Indication</i>	Social, Post-dates, Elective >40 weeks, Elective, “Other”, Multiple gestation, Fetal anomalies
<i>Anticipated Operative Delivery Indication</i>	History of prior uterine surgery, Multiple gestation, Malpresentation, Elective