

Table 1. Relationship between postpartum pain type and opioid dose consumption in hospital.

Type of Pain	Oxycodone Dose (unadjusted)		Oxycodone Dose (adjusted)	
	Estimate	<i>P</i> -value	Estimate	<i>P</i> -value
Affective/Evaluative	1.6282	< 0.0001	1.0221	0.0005
Visceral/Nociceptive	0.01692	0.7809	0.2392	0.0003
Somatic/Nociceptive	0.5383	0.0261	0.6646	0.0109
Dynamic/Evoked	1.113	0.3785	1.5157	0.2565
Neuropathic	0.07083	0.7184	0.1125	0.5729
Nociceptive	-0.1568	0.0071	-0.394	< 0.0001
Nociceptive and/or Neuropathic	-0.1837	0.2094	-0.1852	0.2156

Table 2. Relationship between postpartum pain type and pain intensity/pain scores by 0-10 numeric rating scale.

Type of Pain	Pain Intensity (unadjusted)		Pain Intensity (adjusted)	
	Estimate	<i>P</i> -value	Estimate	<i>P</i> -value
Affective/Evaluative	1.8339	< 0.0001	1.7874	< 0.0001
Visceral/Nociceptive	0.07784	0.0013	0.1078	< 0.0001
Somatic/Nociceptive	0.9289	< 0.0001	0.8918	< 0.0001
Dynamic/Evoked	0.5249	0.296	0.4551	0.4009
Neuropathic	0.7932	< 0.0001	0.7593	< 0.0001
Nociceptive	-0.175	< 0.0001	-0.1972	< 0.0001
Nociceptive and/or Neuropathic	0.05728	0.3169	-0.0030	0.9621